FOR INFORMATION PURPOSES ONLY

The following examples are not meant to be a recommendation or advise for any individual. Only your personal physician can determine your therapy after a complete review of your condition. Consult your physician for specific advise on therapy choices or any questions.

Year 2006

CHEMOTHERAPY OF ADVANCED COLO-RECTAL CANCER (STAGE 4) :EXAMPLES OF AVAILABLE REGIMENS

FU-LV	**C Y C L E 1**								**Avastin can be added every 2 weeks**
NSABP							OFF	OFF	
ROSEWELL PARK	**week 1**	**2**	**3**	**4**	**5**	**6**	**7**	**8**	

ORAL XELODA	**C Y C L E 1**	**C Y C L E 2**	**continue**
	7 days off	**7 days off**	
	days 1-14 **days 14-21 off**	**days 1-14** **days 14-21 off**	

FOLFOX-antibody	**DAY 1**	**DAY 2**	**Day 3**
	FU pump over 22 hours	**FU pump over 22 hours**	Avastin or Erbitux
	Oxal+LV	**LV**	

FOLFIRI-antibody	**DAY 1**	**DAY 2**	**Day 3**
	FU pump over 22 hours	**FU pump over 22 hours**	Avastin or Erbitux
	Iri+LV	**LV**	

Participation in a Clinical trial is always one of the best options.

HOPE CANCER CLINIC, 14555 LEVAN ROAD, SUITE 408E, LIVONIA, MI 48154. PH:734-462-2990, FAX: 734-462-3268.
PREPARED BY HARMESH NAIK, MD.